

TUBERCULOSIS RISK ASSESSMENT FORM

Physician/ Health Provider:		Phone:	Phone:		Date:		
Name:			Date	of Birth:	//_		
Address:				e:	County:		
Phone:	Race: 🛛 White 🗆 Black 🗆 Asian 🗗 Am. Indian/Nat. Alaskan 🗇 Other						
Sex: DMale DFemale	Hispanic: 🗆 No 🗇 Yes	US Born: 🗗 Yes 🗖 No	lf no, US Da	ite of Arri	val://_		

TB RISK FACTORS AND MEDICAL CONDITIONS:

1. Have you ever had a positive skin or blood test?	□Yes □No	If yes, when:
2. Have you had any symptoms of TB (cough, fever, night sweats, loss of appetite, weight loss, or fatigue)?	□Yes □No	If yes, name of symptoms:
3. Have you ever been told you have an abnormal chest X-Ray?	□Yes □No	If yes, when:
4. In the last 2 years, have you lived with or spent time with someone who has been sick with TB?	□Yes □No	
5. Were you born in Africa, Asia, Pacific Islands (except Japan), Central America, South America, Mexico, Eastern Europe, The Caribbean, or the Middle East?	□Yes □No	If yes, in what country were you born:
6. Have you lived or traveled in Africa, Asia, Pacific Islands (except Japan), Central America, South America, Mexico, Eastern Europe, The Caribbean, or the Middle East for more than one month?	□Yes □No	If yes, in what country did you trave to:
7. Have any members of your household come to the United States from another country?	□Yes □No	If yes, name of country:
 8. Have you been exposed to a person who: Is currently in jail or who has been in jail in the past 5 years? Has HIV? Is homeless? Lives in a group home? Uses illegal drugs? Is a migrant farm worker? 	□Yes □No	If yes, name the risk factors you have been exposed to:
 9. Do you have any of the following medical conditions? Diabetes Chronic kidney failure with dialysis Cancer of the neck, head, or lungs Cancer of the blood or lymph system HIV/AIDS Autoimmune disease or immunosuppressive condition Intestinal bypass or gastrectomy 	□Yes □No	
10. Have you ever been in jail or prison?	□Yes □No	If yes, where:
11. Have you ever been an injection drug user?	□Yes □No	If yes, specify:
12. Have you had an organ transplant?	□Yes □No	If yes, specify:
13. Have you been around a person sick with active TB disease?	□Yes □No	If yes, when:
14. Have you ever worked in a lab that processed TB samples?	□Yes □No	If yes, where:

If you answered NO to all of the above questions, you are not in a high-risk group and do not need a TB skin test.

If you answered YES to any of the above questions, you fall into a high-risk group and should have a TB skin test or other tests for TB.

MEDICAL INFORMATION:					
Primary Reason for Evaluation: Contact Investigation					
Dother:					
Symptomatic: INO IYes If Yes, ONSET date:/					
Symptoms: Cough CHemoptysis Fever Night Sweats Weight Loss oflbs					
Previous BCG Vaccine: DNO DYes					
Tuberculin Skin Test (TST/Mantoux/PPD)	Induration: mm				
Date Given:/	Impression: Negative Positive 				
Date Read://					
Interferon Gamma Release Assay (IGRA)	Impression: Negative Positive Indeterminate 				
Date://					
Chest X-ray (required with positive TST or IGRA)	Impression: Normal Abnormal findings				
Date://					
ITBI treatment (Rx and start date):	Prior TB/LTBI treatment (Rx and duration):				
Rx: Date://	Rx:mm				
Contraindications to INH or rifampin for LTBI	Offered but refused LTBI treatment				

ADDITIONAL COMMENTS:

RECOMMENDATIONS:



For the following persons who are at highest risk of developing active tuberculosis disease if they are infected, tuberculin skin tests are considered **positive at 5mm of induration or larger.**

- Recent contacts of TB case patients
- Persons with fibrotic changes on chest radiograph consistent with prior TB
- Patients with organ transplants and other immunosuppressed patients (e.g., taking the equivalent of >15 mg/day of prednisone for 1month or longer, or taking tumor necrosis factor inhibitors)
- Human Immunodeficiency Virus (HIV/AIDS) infected persons

For the following persons with other known **high-risk medical conditions** or **population risks** for recent infection, tuberculin skin tests are positive at **10mm of induration or larger:**

- Foreign-born persons, including children, from areas that have a high TB incidence or prevalence (e.g. Latin America and the Caribbean, Africa, Asia, Eastern Europe, and Russia)*
- Illicit drug users
- Residents and employees of the following high-risk congregate settings:
 - Correctional institutions
 - Long-term facilities
 - Mental health institutions
 - Child care facilities
 - o Hospitals and other health care facilities
 - o Residential facilities for patients with acquired immunodeficiency syndrome (AIDS)
 - Homeless shelters
- Mycobacteriology laboratory personnel
- Persons with the following clinical conditions that place them at high risk:
 - o Silicosis
 - o Diabetes mellitus
 - Chronic renal failure/end-stage renal disease
 - Hematologic/reticuloendothelial disease
 - Cancer of head, neck, or lung
 - o Intestinal bypass or Gastrectomy
 - $\circ \quad \text{Chronic malabsorption syndromes}$
 - Organ Transplant
- Infants, children, and adolescents frequently exposed to adults at high risk for TB infection
- Children <4 years of age (unless the child falls under another category due to condition or exposure)
- Travelers who have reported history of prolonged exposure or extended stay in endemic county (e.g. Latin America and the Caribbean, Africa, Asia, Eastern Europe, and Russia)

For persons at low risk for TB, for whom tuberculin testing is not generally indicated, tuberculin skin tests are **positive at** 15mm of induration or larger:

• Persons with no known risk factors for TB.

*Additional instructions/recommendations may apply to individuals seeking care through a Civil Surgeon. Please reference recommendations: <u>http://www.cdc.gov/ncidod/dq/civil.htm</u>